



Topology and its Applications 56 (1994) 159–164

**TOPOLOGY
AND ITS
APPLICATIONS**

Refinements of T_1 , compact and second countable topologies *

M. Morayne **

Mathematical Institute, Polish Academy of Sciences, ul. Kopernika 18, Wrocław, Poland

C. Ryll-Nardzewski ***

Institute of Mathematics, Wrocław Technical University, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland

(Received 12 November 1992)

Abstract

We prove that every T_1 , compact and second countable topology has a refinement which is a topology of a Polish space and the Borel sets of the two topologies are the same. This shows immediately that every T_1 , compact and second countable space has the standard Borel structure. A representation of all T_1 , compact and second countable topologies as some special topologies on $P(\mathbb{N})$ is also given.

Key words: Borel sets; T_1 topological spaces; Compact spaces; Separation axioms

AMS (MOS) Subj. Class.: 54H05

We shall use the following notation. \mathbb{N} stands for the set of all positive integers, $P(\mathbb{N})$ denotes the family of all subsets of \mathbb{N} . Assigning to each $A \in P(\mathbb{N})$ the sequence $(i_n; n \in \mathbb{N})$ such that $i_n = 1$ if $n \in A$ and $i_n = 0$ if $n \notin A$ we shall consider $P(\mathbb{N})$ as $\{0, 1\}^{\mathbb{N}}$ with the product topology. If \mathcal{U} is any family of subsets of a set X , then by $\mathcal{O}(\mathcal{U})$ we shall denote the topology generated by the subbase \mathcal{U} . A *refinement* of a topology \mathcal{O} is any topology richer than \mathcal{O} . For A being a subset of

* This paper was supported in part by KBN Grant 2-1054-91-01.

** This paper was written in part while the author was a postdoctoral fellow at the Department of Mathematics, York University, Toronto, Canada. At that time the author was supported from NSERC grants of Professor Alan Dow, Professor Claude Laflamme and Professor Juris Steprāns.

*** Corresponding author.

a product of sets by $\pi_n(A)$ we shall denote the projection of A onto the n th axis. The symbol $\text{diag}(X^2)$ will stand for the set $\{(x, x) : x \in X\}$. For a topological space (X, \mathcal{O}) and $A \subseteq X$ by $\mathcal{O}|A$ we shall denote the relative topology \mathcal{O} on A .

A family \mathcal{A} of sets is *compact in the sense of Marczewski* if for every $\mathcal{A}' \subseteq \mathcal{A}$ there is $\bigcap \mathcal{A}' \neq \emptyset$ whenever $\bigcap \mathcal{A}'' \neq \emptyset$ for all finite $\mathcal{A}'' \subseteq \mathcal{A}'$. If \mathcal{A} is any family of sets then \mathcal{A}_s will denote the family of all finite unions of elements of \mathcal{A} .

It is known that if A and B are Borel subsets of Polish spaces and A and B are of the same cardinality then the σ -algebras of Borel subsets of A and B are isomorphic. We say that a topological space X has a *standard Borel structure* if the σ -algebra of Borel subsets of X is isomorphic to the σ -algebra of Borel subsets of a Polish space (in [2] the term *standard Borel space* refers to a pair consisting of a set X and a σ -algebra of its subsets which contains X and which is isomorphic to the σ -algebra of Borel subsets of a Polish space, and X is not endowed with a topology).

We start with characterizing T_1 compact and second countable topological spaces. We shall extensively use the idea of Marczewski's characteristic function of sequence of sets (first used by Marczewski (Szpilrajn) in [6]).

Let \mathcal{J} be an arbitrary nonempty family of finite subsets of \mathbb{N} which is hereditary with respect to inclusion, i.e., satisfying

$$(J_1 \subseteq J_2 \text{ and } J_2 \in \mathcal{J}) \text{ implies } J_1 \in \mathcal{J}. \quad (1)$$

Now we define two families $\mathcal{G}, \mathcal{H} \subseteq P(\mathbb{N})$. The former consists of all sets the finite subsets of which are in \mathcal{J} , the latter consists of those elements of \mathcal{G} which are maximal with respect to inclusion:

$$\mathcal{G} = \{G \in P(\mathbb{N}) : (\forall n \in \mathbb{N}) [[1, n] \cap G \in \mathcal{J}]\} \quad (2)$$

and

$$\mathcal{H} = \{H \in \mathcal{G} : (\forall G \in \mathcal{G}) [H \subseteq G \rightarrow H = G]\}. \quad (3)$$

Note that \mathcal{G} is a closed subset of $P(\mathbb{N})$; in fact it is easy to observe that \mathcal{G} is the closure of \mathcal{J} (recall that $P(\mathbb{N})$ is identified with $\{0, 1\}^{\mathbb{N}}$ with the product topology).

Let us now consider the family

$$\mathbf{K} = \{\mathcal{K}_n : n \in \mathbb{N}\}, \quad (4)$$

where

$$\mathcal{K}_n = \{H \in \mathcal{H} : n \in H\}. \quad (5)$$

We shall show that \mathbf{K} is compact in the sense of Marczewski. Let $\mathbf{K}' = \{\mathcal{K}_{n_1}, \mathcal{K}_{n_2}, \dots\} \subseteq \mathbf{K}$, and let us assume that $\mathcal{K}_{n_1} \cap \dots \cap \mathcal{K}_{n_m} \neq \emptyset$, for each $m \in \mathbb{N}$. We have to show that $\bigcap \mathbf{K}' \neq \emptyset$. Pick an element $H_m \in \mathcal{K}_{n_1} \cap \dots \cap \mathcal{K}_{n_m}$. As \mathcal{G} is a closed subset of $P(\mathbb{N})$ we can find a subsequence $(H_{m_i})_i$ which is convergent to some $G \in \mathcal{G}$. Take any $H \in \mathcal{H}$ such that $G \subseteq H$. It is easy to observe that the set $\{n_1, n_2, \dots\}$ is contained in H and thus $H \in \bigcap \mathbf{K}'$. We have proved that \mathbf{K} is compact in the sense of Marczewski, whence, as it is well known (see for instance [5, III, 1, the proof of Theorem 4]), \mathbf{K}_s has the same property. Now we can

consider \mathcal{H} as a topological space taking $K^c = \{\mathcal{H} \setminus \mathcal{K} : \mathcal{K} \in K\}$ as a subbase. Obviously this space is second countable and compact. It is also T_1 because if $H_1, H_2 \in \mathcal{H}$ and $H_1 \neq H_2$, then as H_1 is maximal, there is $n \in H_1 \setminus H_2$, whence $H_1 \in \mathcal{H}_n$ and $H_2 \notin \mathcal{H}_n$.

We shall prove now that the construction above describes all possibilities. More exactly, the following theorem holds.

Theorem 1. *Every T_1 , compact and second countable topological space X is homeomorphic to $(\mathcal{H}, \mathcal{O}(K^c))$ for some family \mathcal{J} of finite subsets of \mathbb{N} satisfying (1), where \mathcal{H} and K are defined by (2), (3), (4) and (5).*

Proof. Let \mathcal{U} be a countable base for X and let $\mathcal{C} = \{C_1, C_2, \dots\} = \mathcal{U}^c = \{X \setminus U : U \in \mathcal{U}\}$. Let $\mathcal{J} = \{J : J \text{ is a finite subset of } \mathbb{N} \text{ such that } \bigcap \{C_n : n \in J\} \neq \emptyset\}$. The families \mathcal{G} and \mathcal{H} are defined from \mathcal{J} as previously (see (2) and (3)). Let $x \in X$ and $\phi(x) = \{n : x \in C_n\}$. We shall show now that ϕ is a homeomorphism between X and \mathcal{H} . We have $\bigcap \{C_n : n \in \phi(x)\} = \{x\}$ because X is T_1 , and thus $\phi(x) \in \mathcal{H}$. Now let $H \in \mathcal{H}$. By the definition of \mathcal{G} we have $\bigcap \{C_n : n \in A\} \neq \emptyset$ for any finite subset A of H , whence $\bigcap \{C_n : n \in H\} \neq \emptyset$ because X is compact. If $x_1, x_2 \in \bigcap \{C_n : n \in H\}$ and $x_1 \neq x_2$, then there is $C_n \in \mathcal{C}$ such that $x_1 \in C_n$ and $x_2 \notin C_n$ because X is T_1 . But then $H \cup \{n\} \in \mathcal{G}$ which is a contradiction with the definition of \mathcal{H} . Hence $\bigcap \{C_n : n \in H\} = \{x\}$ for some $x \in X$ and, obviously, $\phi(x) = H$. Thus ϕ is a 1–1 mapping from X onto \mathcal{H} . To prove that ϕ is a homeomorphism it is enough to notice that $\phi(C_n) = \mathcal{H}_n$. \square

We shall use the above characterization of T_1 compact and second countable topologies to prove that for each such topology there exists a finer one which is a topology of a Polish space generating the same Borel sets. Namely we shall prove the following theorem.

Theorem 2. *If (X, \mathcal{O}) is a T_1 , compact and second countable topological space, then there is a topology \mathcal{O}^* which is a refinement of \mathcal{O} such that (X, \mathcal{O}^*) is metrizable as a Polish space and the Borel sets generated by \mathcal{O}^* are the same as those generated by \mathcal{O} .*

In the proof of the above theorem we shall use the following fact which we shall prove first.

Fact 3. *If X is a metrizable compact space, R is a binary relation on X and R is a closed subset of X^2 then the set of all elements maximal in R is a G_δ subset of X .*

Proof. It is enough to notice that

$$\{x : x \text{ is maximal with respect to } R\} = \left(\pi_1(R \setminus \text{diag}(X^2)) \right)^c. \quad \square$$

Proof of Theorem 2. In the view of Theorem 1 it is enough to prove the theorem for each space of the form $(\mathcal{H}, \mathcal{O}(K^c))$ as in Theorem 1.

We have

$$\mathcal{H}_n = \{A \in P(\mathbb{N}) : n \in A\} \cap \mathcal{H},$$

whence \mathcal{H}_n is a clopen subset of \mathcal{H} with the relative topology inherited from $P(\mathbb{N})$. Hence this topology extends that generated by a subbase K^c .

The sets $\{A \in P(\mathbb{N}) : n \in A\}$ are generators of the σ -algebra of Borel sets in $P(\mathbb{N})$ and \mathcal{H}_n are closed sets in \mathcal{H} with the topology $\mathcal{O}(K^c)$. Thus \mathcal{H} with the relative topology inherited from $P(\mathbb{N})$ has the same Borel sets as with the topology $\mathcal{O}(K^c)$.

Thus the proof is complete because, as it follows from Fact 3, \mathcal{H} is a G_δ subset of a Polish space $P(\mathbb{N})$ and thus is itself metrizable as a Polish space. \square

The following corollary is immediate.

Corollary. Every T_1 , compact and second countable space has standard Borel structure.

The following example due to Dow [3] shows that one cannot always obtain a Hausdorff and compact refinement of the T_1 , compact and second countable topology.

Example 4 (Dow). Let $a, b \notin (0, 1]$, $a \neq b$, and

$$X = (0, 1] \cup \{a, b\}.$$

Let us define the neighbourhood base for a to consist of all sets of the form

$$U_n = \{a\} \cup \bigcup \left\{ ((2i+2)^{-1}, (2i)^{-1}) : i \geq n \right\},$$

$n \in \mathbb{N}$, and the neighbourhood base for b to consist of all sets of the form

$$V_n = \{b\} \cup \bigcup \left\{ ((2i+1)^{-1}, (2i-1)^{-1}) : i \geq n \right\},$$

$n \in \mathbb{N}$. The points of $(0, 1]$ have their usual neighbourhood bases. As any compact and Hausdorff refinement of this topology must remain the same on $(0, 1]$ it would be a two-point compactification of $(0, 1]$ and such compactification does not exist for the remainder of any compactification of $(0, 1]$ must be connected (see, for instance, [1, the first lines of the proof of Theorem 1]).

Below we give an example of a T_1 , compact and second countable topology \mathcal{S} that does not have any Hausdorff, σ -compact refinement consisting of Borel sets of \mathcal{S} . First we state the following fact.

Fact 5. The space X described in Example 4 has the property that if K is its compact and dense subset, then $K = X$.

Proof. It is easy to see that $K \cap (0, 1]$ must be dense in $(0, 1]$, and thus $(0, 1] \subseteq K$. If K did not contain either of the points a, b , then it would not be compact. The case where exactly one of the points a, b lies in K is also impossible. In fact if, for instance, $K = (0, 1] \cup \{a\}$, then the open cover of K consisting of the intervals $((2i+1)^{-1}, (2i-1)^{-1})$, $i \geq 1$, the set U_1 defined in Example 4 and the interval $(2^{-1}, 1]$ does not have any finite subcover, which contradicts the compactness of K . Thus we have $K = X$. \square

Example 6. Let (X, \mathcal{O}) be a T_1 , compact and second countable topological space having no Hausdorff and compact refinement. Assume also that (X, \mathcal{O}) has the property that each set compact and dense in X must be the whole X . The space described in Example 4 has all the properties required (see Fact 5).

Let the space $Y = X^\omega$ be equipped with the usual product topology \mathcal{S} . Of course, (Y, \mathcal{S}) is T_1 , compact and second countable. We claim that this space does not have any Hausdorff, σ -compact refinement consisting of Borel sets of \mathcal{S} .

Let \mathcal{O}^* be a refinement of \mathcal{O} as in the conclusion of Theorem 2.

Let \mathcal{P} be the topology on Y defined as a product of countably many copies of \mathcal{O}^* .

Assume now that \mathcal{V} is a Hausdorff refinement of \mathcal{S} consisting of Borel sets generated by \mathcal{S} . To prove that \mathcal{V} cannot be σ -compact it is enough to show that every set $K \subseteq Y$ compact in the topology \mathcal{V} is of the first category in the topology \mathcal{P} (for (Y, \mathcal{P}) is metrizable as a Polish space).

Assume for a contradiction that K is of second category in the topology \mathcal{P} . The set K is closed in the topology \mathcal{V} , hence it is a Borel set in \mathcal{P} , thus possessing Baire's property in \mathcal{P} . By [4, Chapter 1, 11, IV, Corollary 2] there exist sets U_1, \dots, U_n open in \mathcal{O}^* such that the set $K \cap (U_1 \times \dots \times U_n \times X \times X \times \dots)$ is residual in the set $U_1 \times \dots \times U_n \times X \times X \times \dots$ (in the topology \mathcal{P}). By the Kuratowski–Ulam theorem [4, Chapter 2, 22, V, Corollary 1a] there is a sequence of points $c_1 \in U_1, \dots, c_n \in U_n, c_{n+2} \in X, c_{n+3} \in X, \dots$ such that the set

$$S = \{x \in X : (c_1, \dots, c_n, x, c_{n+2}, c_{n+3}, \dots) \in K\}$$

is residual and, therefore, dense in X with the topology \mathcal{O}^* . This implies that S is also dense in X in the topology \mathcal{O} , because \mathcal{O}^* is a refinement of \mathcal{O} .

$$T = \{(c_1, \dots, c_n, x, c_{n+2}, c_{n+3}, \dots) : x \in X\}$$

is closed in X^ω with the topology \mathcal{S} , whence it is also closed in the topology \mathcal{V} . Hence the set $T \cap K$ is closed in K with the topology $\mathcal{V}|_K$ and, because K is compact in this topology, $T \cap K$ is compact in the topology \mathcal{V} . Thus the set $T \cap K$ is also compact in the topology \mathcal{S} , whence its projection $S = \pi_{n+1}(T \cap K)$ is compact in the topology \mathcal{O} . So now we know that S is both dense and compact in X in the topology \mathcal{O} , whence $S = X$. Thus the projection of the topology $\mathcal{V}|_{T \cap K}$ (note that $T \cap K = T$) onto the $n+1$ axis would be a Hausdorff and compact refinement of the topology \mathcal{O} , but such a refinement does not exist.

Remark. It follows from Theorem 2 that any T_1 , compact and second countable topology has a refinement which is a topology of a G_δ subset of a Hausdorff, compact and second countable space. Using Example 6 we can show that this result cannot be improved, namely that there is a T_1 , compact and second countable topology that does not have a refinement which is a topology of an F_σ subset of a Hausdorff, compact and second countable topological space. In particular, it will not have a Hausdorff, locally compact and second countable refinement.

The space (Y, \mathcal{S}) of Example 6 is just the one we need. Assume for a contradiction that it has a refinement which is the topology \mathcal{T} of an F_σ subset of a Hausdorff, compact and second countable topological space. By Theorem 2 the space (Y, \mathcal{S}) has the standard Borel structure, namely that of the space (Y, \mathcal{P}) of Example 6. But then the identity function $\text{id}: Y \rightarrow Y$, where the domain is equipped with the topology \mathcal{P} and the range with the topology \mathcal{T} , is a Borel homeomorphism of Y [4, Chapter 3, 39, V, Theorem 1] and all Borel sets of \mathcal{T} coincide with those of \mathcal{P} , and therefore of \mathcal{S} . Thus the refinement in question would consist of Borel sets of \mathcal{S} , but it does not exist as it was shown in Example 6.

1. Acknowledgement

We are grateful to Professors Alan Dow, Claude Laflamme, Juris Steprāns and Steve Watson for their comments concerning this paper and to them as well as to the other members of York University topology seminar for their nice and stimulating interest in our problems.

2. References

- [1] D.P. Bellamy, A non-metric indecomposable continuum, *Duke Math. J.* 38 (1971) 15–20.
- [2] K.P.S. Bhaskara-Rao and B.V. Rao, Borel spaces, *Dissertationes Math.* 190 (1981).
- [3] A. Dow, Private communication to the authors.
- [4] K. Kuratowski, *Topology I* (Academic Press, New York, 1966).
- [5] P.A. Meyer, *Probability and Potentials* (Blaisdell Publishing Company, Waltham, MA, 1966).
- [6] E. Szpilrajn, Sur l'équivalence des suites d'ensembles et l'équivalence des fonctions, *Fund. Math.* 26 (1936) 302–326.